

Polaronic Quasiparticles in a Strongly Correlated Electron Band

W. Koller,^{*} A. C. Hewson,[†] and D. M. Edwards[‡]

Department of Mathematics, Imperial College, London SW7 2AZ, UK

(Dated: 18 July 2005)

We show that a strongly renormalized band of polaronic quasiparticle excitations is induced at the Fermi level of an interacting many-electron system on increasing the coupling of the electrons to local phonons. We give results for the local density of states at zero temperature both for the electrons and phonons. The polaronic quasiparticles satisfy Luttinger's theorem for all regimes considered, and their dispersion shows a kink similar to that observed experimentally in copper oxides. We calculate the quasiparticle weight factor z and deduce the local effective inter-quasiparticle interaction $\tilde{U}$. Our calculations are based on the dynamical mean field theory and the numerical renormalization group for the hole-doped Holstein-Hubbard model and large on-site repulsion.

PACS numbers: 71.10.Fd, 71.30.+h, 71.38.-k

I. INTRODUCTION

Polarons as quasiparticles, in which the electronic excitations are strongly coupled to the lattice modes, have been studied theoretically for well over 50 years. Nearly all these studies have been confined to models with only one or two electrons, and do not include the electron spin. Such models have limited applicability and there is a need for studies of more realistic many-electron situations¹. There are technical problems, however, in handling such models as non-perturbative methods are required to describe the effects of strong electron-lattice interactions and few such methods are applicable to the many-electron case. Progress has been made in recent years in dealing with models with strong local interactions via the dynamical mean field theory (DMFT)^{2,3}. Calculations based on the Holstein and Holstein-Hubbard models, where the electrons are coupled to local Einstein lattice modes, have been made using the DMFT to examine the effect of strong electron-phonon interactions on the Mott transition^{4,5}, for example. The Holstein model⁶, however, was originally introduced to describe small polaron effects. Within DMFT, only the single electron case can be solved exactly^{7,8}. There have been few studies at finite electron densities and including the electron spin⁹, except in one dimension^{10,11}. The problem is that if spin is included in the Holstein model, there is an effective attractive local interaction between the electrons, which leads to local bipolaron formation and, away from half filling, superconductivity.

Once the electron spin is included the more realistic model is the Holstein-Hubbard model which includes an local electron repulsion U . In most situations the repulsion U dominates and undermines any tendency to the formation of local bipolarons. In the Holstein-Hubbard model at half filling with a large U , the electron-phonon interaction plays no important role as charge fluctuations are suppressed; its main effect is to renormalise and reduce the effective value of U ^{4,5}. At half filling and small U , where the electron-phonon coupling dominates, again there is little evidence⁴ of purely polaronic effects, because of the formation of bipolarons^{4,12}. The question

arises, therefore, whether there is any parameter regime where purely polaronic effects can be observed. If there is, then does the behaviour correspond to the predictions based on the spinless models with only one or two electrons, or do many-electron correlations play a significant role?

In this letter we present calculations to show clearly that small strongly renormalized polaronic excitations do exist in the Holstein-Hubbard model and form a distinct narrow band at the Fermi level. The Hamiltonian for this model is

$$H = \sum_{\mathbf{k}\sigma} \epsilon(\mathbf{k}) c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma} + U \sum_i n_{i\uparrow} n_{i\downarrow} + \omega_0 \sum_i b_i^\dagger b_i + g \sum_i (b_i^\dagger + b_i)(n_{i\uparrow} + n_{i\downarrow} - 1), \quad (1)$$

where $\epsilon(\mathbf{k})$ describes the dispersion of the band electrons and U is the on-site inter-electron repulsion as in the Hubbard model¹³. The electron density $n_i = n_{i\uparrow} + n_{i\downarrow}$ at site i couples linearly to the local displacement operator $x_i \equiv (b_i + b_i^\dagger)/\sqrt{2m\omega_0}$ with an electron-phonon coupling g . The phonons are assumed to be dispersionless (local Einstein phonons) with energy ω_0 and m is the mass of the vibrating ions.

Our calculations are based on DMFT using a semi-elliptical density of states $\rho_0(\omega)$ of the non-interacting system given by

$$\rho_0(\omega) = \frac{2}{\pi D^2} \sqrt{D^2 - (\omega + \mu_0)^2} \quad (2)$$

for $|\omega + \mu_0| \leq D$ and zero elsewhere, corresponding to a Bethe lattice, where $2D$ is the bandwidth and μ_0 is the chemical potential. The effective impurity problem is solved using the numerical renormalization group (NRG)^{14,15}, as generalized for the calculation of dynamical response functions^{16,17,18}, and incorporating the density matrix generalization¹⁹ due to Hofstetter.

To suppress any tendency for bipolaron formation we take a large value of the on-site repulsion, $U = 6$ in units where $D = 2$, which for half filling corresponds to a Mott insulator^{4,20}. We choose a value of μ corresponding to




FIG. 1: Local density of states for the quarter-filled Hubbard model ($U = 6.0$)

quarter filling $\langle n_i \rangle = 0.5$. The local density of states for $g = 0$ is shown in Fig. 1, with the Fermi level such that the lower Hubbard band is $1/3$ filled. In this regime the strong correlation effects are largely suppressed due to the charge fluctuations; there is no sharp quasiparticle peak at the Fermi level, which occurs for the small doping regime near half filling, and there is a moderately reduced quasiparticle renormalization factor $z \approx 0.65$. The small spectral weight between the upper and lower bands is due to the tails in the broadening in the higher energy peaks of the discrete NRG results.

To study the impact of electrons coupling to phonons we concentrate on the lower Hubbard band. In Fig. 2 we show the effect of increasing the electron-phonon interaction g , taking $\omega_0 = 0.2$ in all cases. As g increases to the value $g = 0.65$ a sharp feature develops at the Fermi level which we attribute to a polaronic renormalization. Due to the large U , there is negligible double occupancy $\langle n_{i\uparrow}n_{i\downarrow} \rangle < 0.01$ for this range of g indicating no local bipolaron formation.

To look at the band aspects of these excitations, we introduce the $\mathbf{k}$ -resolved one-electron Green's function $G_{\mathbf{k},\sigma}(\omega)$ with a local self-energy $\Sigma_\sigma(\omega)$,

$$G_{\mathbf{k},\sigma}(\omega) = \frac{1}{\omega + \mu - \epsilon(\mathbf{k}) - \Sigma_\sigma(\omega)}. \quad (3)$$

Figure 3 shows a plot of the spectral density of the Green's function $G_{\mathbf{k}}(\omega)$ for $-2 \leq \epsilon(\mathbf{k}) \leq 2$ and $g = 0.60$. When integrated over $\mathbf{k}$, it gives the local density of states for the corresponding g shown in Fig. 2. We plot the positions of the maxima in the quasiparticle peaks as a function of $\epsilon(\mathbf{k})$ in Fig. 4. The flat dispersion in the region of the Fermi level is clear evidence of a narrow polaronic band (full line). We see a kink at $\omega \approx -0.05$ which we discuss later in relation with the phonon spectra. None of this is seen in the pure Hubbard case, shown as the dashed line in Fig. 4.

The locus of points $\mathbf{k}_F$ on the Fermi surface of the interacting system is given by

$$\epsilon(\mathbf{k}_F) = \mu - \Sigma(0). \quad (4)$$




FIG. 2: The development of the polaronic quasiparticle band in the lower Hubbard band of the quarter-filled Holstein-Hubbard model ($U = 6.0$) for increasing electron-phonon coupling g .




FIG. 3: Spectral density $\rho_k(\omega)$ of the electron Green's function for the quarter-filled Holstein-Hubbard model ($U = 6.0, g = 0.60$).

According to Luttinger's theorem²¹ the volume of the Fermi surface should be the same as that of the non-interacting system. For a local self energy $\Sigma(\omega)$, this implies that the Fermi surface given by Eq. (4) should be the same as the non-interacting one, $\epsilon(\mathbf{k}_F) = \mu_0$. If these are to coincide, then $\mu_0 = \mu - \Sigma(0)$.




FIG. 4: Positions of the maxima of the spectral function $\rho_k(\omega)$ of the electron Green's function for the quarter-filled Holstein-Hubbard model ($U = 6.0, g = 0.60$, full line) and the quarter-filled Hubbard model ($U = 6.0, g = 0.00$, dashed line). The arrow indicates $\epsilon(k_F)$.

To check the theorem we have calculated the density $\langle n_i \rangle$ directly from the NRG expectation value in the ground state, and then from the non-interacting Fermi surface using the value of $\mu_0 = \mu - \Sigma(0)$ from equation (4). The two sets of results are compared for a range of hole dopings in Fig. 5 (left) for the pure Hubbard model, and in Fig. 5 (right) for the Holstein-Hubbard model with $g = 0.5$. The agreement between the two sets of results provide a convincing evidence of the applicability of Luttinger's theorem in strongly correlated systems both with and without the electron-phonon interaction.

If we assume Fermi liquid theory²¹ and expand the self-energy in Eq. (3) about the Fermi level $\omega = 0$, and retain the first two terms, we can define a quasiparticle Green's function

$$\tilde{G}_{\mathbf{k},\sigma}(\omega) = \frac{1}{\omega + \tilde{\mu} - \tilde{\epsilon}(\mathbf{k})} \quad (5)$$

where $\tilde{\epsilon}(\mathbf{k}) = z\epsilon(\mathbf{k})$, and z is the usual wavefunction renormalization factor $z = (1 - \Sigma'(0))^{-1}$, and $\tilde{\mu} = z(\mu - \Sigma(0))$ is a renormalized chemical potential. The corresponding density of states $\tilde{\rho}_0(\omega)$ for the non-interacting quasiparticles is given by

$$\tilde{\rho}_0(\omega) = \frac{2}{\pi \tilde{D}^2} \sqrt{\tilde{D}^2 - (\omega + \tilde{\mu})^2} \quad (6)$$

where $\tilde{D} = zD$ plays the role of a renormalized band width. We interpret this as a renormalized band of free polaronic quasiparticles. Integrating this free quasiparticle band up to the Fermi level gives the quasiparticle occupation $\langle \tilde{n}_i \rangle$. The resulting $\langle \tilde{n}_i \rangle = \langle n_i \rangle$, demonstrating the one-to-one correspondence of the electron and quasiparticle states of Fermi liquid theory^{22,23} (see Fig. 5).

In Fig. 6 we give the results for the change in the quasiparticle renormalization factor z with increase of g for quarter filling and $U = 6$. The values of z we calculated by differentiating the self-energy, and also from an




FIG. 5: Density as a function of the chemical potential μ calculated directly (full lines) and from the volume of the Fermi surface (circles) for the Hubbard model (left-hand plot, $U = 6$) and the Holstein-Hubbard model (right-hand plot, $U = 6, g = 0.5$).




FIG. 6: Quasiparticle weight as a function of g for the quarter-filled Holstein-Hubbard model ($U = 6.0$) deduced from the derivative of the self energy (o) and from an analysis of the renormalized parameters (x).

analysis of the renormalized parameters^{24,25} on the approach to the low energy fixed point. The two types of estimates are in reasonable agreement; the deviation for smaller values of g is largely due to fluctuations in the numerical differentiation of $\Sigma(\omega)$ in this range. The behaviour of z with increase of g is quite different from that near half filling where z increases with increase of g . Near half filling the strong correlation effects stem from the enhancement of spin fluctuations and the suppression of charge fluctuations due to the large value of U . In this regime the increase of z with g can be understood as due to the reduction in the effective value of U , $U_{\text{eff}} = U - 2g^2/\omega_0$; hence an increase in g reduces U_{eff} and hence reduces the renormalization^{4,5}. The results for z at quarter filling shown in Fig. 6, by contrast, show a large decrease of z with increase of g . This increased renormalization is a pure polaronic effect; the coupling to the lattice causing an increase in the effective mass of the conduction electrons and a reduced mobility.

The renormalized parameter approach^{24,25} also allows us to calculate the local interaction between the quasiparticles $\tilde{U}$. The results for $\tilde{U}$ over this range are shown in Fig. 7. There is a progressive reduction with increase of g as might be expected for the induced attractive interaction due to phonon exchange. For the largest




FIG. 7: Effective local interaction $\tilde{U}$ between quasiparticles as a function of g for the quarter-filled Holstein-Hubbard model ($U = 6.0$).




FIG. 8: Spectral density of the phonon propagator for the quarter-filled Holstein-Hubbard model ($U = 6.0$) for various values of the electron-phonon coupling.

value of g the value of $\tilde{U}$ becomes negative, even though $U - 2g^2/\omega_0 > 0$. This can be interpreted as due to the fact the very low energy quasiparticles hardly ever occupy the same site due to correlation effects (double oc-

cupancy < 0.01) and their effective attraction is due to retardation effects.

For values of $g > 0.65$ the dynamical mean field theory no longer converges satisfactorily, the iterations fluctuate between occupied and unoccupied states. This indicates the on-set of a broken symmetry state, probably some form of charge ordering or phase separation. This instability is also revealed in the one-phonon propagator $d(\omega) = \langle\langle b_i; b_i^\dagger \rangle\rangle_\omega$ whose spectral density is shown in Fig. 8. With increasing g there is a progressive softening of the phonon mode from its bare value $\omega_0 = 0.2$. For $g \approx 0.65$ considerable spectral weight develops for $\omega < 0$. This weight relates to the expected number of excited phonons in the ground state. It clearly indicates an incipient lattice instability for $g > 0.65$.

The kink to be seen at $\omega = -0.05$ in the quasiparticle dispersion in Fig. 4 is clearly related to the frequency of the renormalized phonon excitation at the corresponding $g = 0.6$ (Fig. 8). An analysis for different values of g shows a clear correlation of the kink energy to the *renormalized* phonon frequency. This is similar to the experimentally observed kink in the hole-doped copper oxides²⁶.

To summarize: We have shown that an accurate treatment of the Holstein-Hubbard model at *finite electron density* and $T = 0$ reveals a clear polaronic quasiparticle band induced at the Fermi level. Luttinger's theorem is shown to be satisfied. A kink found in the quasiparticle dispersion relation is related to the renormalized phonon frequency.

Acknowledgments

We wish to thank the EPSRC (Grant GR/S18571/01) for financial support.

-
- * Electronic address: w.koller@imperial.ac.uk
 † Electronic address: a.hewson@imperial.ac.uk
 ‡ Electronic address: d.edwards@imperial.ac.uk
¹ E. V. L. de Mello and J. Ranninger, Phys. Rev. B **58**, 9098 (1998).
² A. Georges, G. Kotliar, W. Krauth, and M. J. Rozenberg, Rev. Mod. Phys. **68**, 13 (1996).
³ M. Jarrell, Phys. Rev. Lett. **69**, 168 (1992).
⁴ W. Koller, D. Meyer, and A. C. Hewson, Phys. Rev. B **70**, 155103 (2004).
⁵ G. Sangiovanni, M. Capone, C. Castellani, and M. Grilli, Phys. Rev. Lett. **94**, 026401 (2005).
⁶ T. Holstein, Ann. Phys. **8**, 325 (1959).
⁷ H. Sumi, J. Phys. Soc. Jpn. **36**, 770 (1974).
⁸ S. Ciuchi, F. de Pasquale, S. Fratini, and D. Feinberg, Phys. Rev. B **56**, 4494 (1997).
⁹ M. Capone and S. Ciuchi, Phys. Rev. Lett. **91**, 186405 (2003).

- ¹⁰ G. Wellein and H. Fehske, Phys. Rev. B **56**, 4513 (1997).
¹¹ H. Fehske, G. Wellein, G. Hager, A. Weiße, and A. R. Bishop, Phys. Rev. B **69**, 165115 (2004).
¹² D. Meyer, A. C. Hewson, and R. Bulla, Phys. Rev. Lett. **89**, 196401 (2002).
¹³ J. Hubbard, Proc. R. Soc. London, Ser. A **276**, 238 (1963).
¹⁴ K. G. Wilson, Rev. Mod. Phys. **47**, 773 (1975).
¹⁵ H. R. Krishna-murthy, J. W. Wilkins, and K. G. Wilson, Phys. Rev. B **21**, 1003 (1980).
¹⁶ O. Sakai, Y. Shimizu, and T. Kasuya, Prog. Theor. Phys. Suppl. **108**, 73 (1992).
¹⁷ T. A. Costi, A. Hewson, and V. Zlatić, J. Phys.: Condens. Matter **6**, 2519 (1994).
¹⁸ R. Bulla, A. C. Hewson, and T. Pruschke, J. Phys.: Condens. Matter **10**, 8365 (1998).
¹⁹ W. Hofstetter, Phys. Rev. Lett. **85**, 1508 (2000).
²⁰ R. Bulla, Phys. Rev. Lett. **83**, 136 (1999).
²¹ J. M. Luttinger, Phys. Rev. **119**, 1153 (1960).

- ²² L. D. Landau, Sov. Phys. JETP **3**, 920 (1957).
- ²³ L. D. Landau, Sov. Phys. JETP **5**, 101 (1957).
- ²⁴ A. C. Hewson, A. Oguri, and D. Meyer, Euro. Phys. J. B. **40**, 177 (2004).
- ²⁵ A. C. Hewson, J. Phys. Soc. Jpn. **74**, 8 (2005).
- ²⁶ A. Lanzara, P. V. Bogdanov, X. J. Zhou, S. A. Keller, D. L. Feng, E. D. Lu, T. Yoshida, H. Eisaki, A. Fujimori, K. Kishio, et al., Nature **412**, 510 (2001).